


**NASA CONTRACTOR
REPORT**



NASA CR-1

2.1

0060409



TECH LIBRARY KAFB, NM

NASA CR-1022

LOAN COPY: RETURN TO
AFWL (WLIL-2)
KIRTLAND AFB, N MEX

MIE-SCATTERING FUNCTION

by F. Shahrokhi and P. Wolf

Prepared by

THE UNIVERSITY OF TENNESSEE SPACE INSTITUTE

Tullahoma, Tenn.

for

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION • WASHINGTON, D. C. • APRIL 1968



0060409

NASA CR-1022

MIE-SCATTERING FUNCTION

By F. Shahrokhi and P. Wolf

Distribution of this report is provided in the interest of information exchange. Responsibility for the contents resides in the author or organization that prepared it.

Prepared under Grant No. NGR-43-001-021 by
THE UNIVERSITY OF TENNESSEE SPACE INSTITUTE
Tullahoma, Tenn.

for

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

MIE-SCATTERING FUNCTION

F. Shahrokhi* and P. Wolf⁺

The University of Tennessee Space Institute
Tullahoma, Tennessee

ABSTRACT

For a comparison between experimental results and Mie-scattering theory, a series of calculations was made to find the normalized scattering function $f(\theta)$ from the tables of Legendre-polynomials by G. C. Clark and S. W. Churchill and from tables of scattering coefficients made by C. M. Chu, G. C. Clark, and S. W. Churchill.

This study was carried out as part of a research program being performed under NASA Sustaining Grant NGR-43-001-021.

*Assistant Professor of Aerospace Engineering.

+ Graduate Research Assistant.

Introduction

The angular distribution of the intensity of radiation scattered at spherical particles was worked out by G. Mie (1) in theory. Several tables were prepared for particle materials of real refractive index (absorption by particle material neglected). In these tables coefficients are given which have to be combined with polynomials to give the scattering distribution function for single scattering at spherical particles of defined size in relation to the radiation wave length. The scattering function is usually for an incident beam of unit intensity.

This combination of coefficients and polynomials was done by a small computer program, based on the data provided by the tables for Legendre polynomials by G. C. Clark and S. W. Churchill (2) and by the tables for scattering coefficients by C. M. Chu, G. C. Clark and S. W. Churchill (3).


The computations became necessary because of an intention to do a closer comparison of experimental results of scattering in particle clouds with scattering predicted by the Mie-theory for spherical particles suspended in a gas.

Mathematical Definitions and Computation

As Hertel (4) proposed, the normalized scattering function $f(\theta)$ may be expressed by

$$f(\theta) = \frac{1}{4\pi} \left(1 + \sum_{n=1}^{\infty} a_n \cdot P_n(\cos \theta) \right)$$

Here is $f(\theta) = \frac{i(\theta)}{\frac{\pi D^2}{4} K}$ the intensity of a beam scattered in direction θ from an incident beam of unit intensity hitting a spherical particle



The scattered intensity $i(\theta)$ is independent of ϕ for spherical particles of uniform material and a uniform irradiation.

K is the scattering cross section nondimensionalized with the geometrical cross section $\frac{\pi}{4} D^2$ of the spherical particle of diameter D .

The scattering coefficients, a_n , are independent of the angle, θ , but vary with particle size referred to radiation wave length $\alpha = \frac{\pi \cdot D}{\lambda}$ in the medium surrounding the particles and with the particle material of refraction relative to the surrounding medium.

The a_n were computed by authors (3) according to the relations derived by them.

$$a_n = \frac{(2)(-1)^n}{\alpha^2 K(\alpha, \beta)} \sum_{j=1}^{\infty} \sum_{k=1}^j \left(\frac{2}{1 + \delta_{jk}} \right) \left\{ (2n+1) \left[\frac{j(j+1) + k(k+1) - n(n+1)}{2} \right]^2 w_{jkn} w_{jk} + v_{jkn} v_{jk} \right\}$$

where

$$\delta_{jk} = \begin{cases} 0 & \text{for } j \neq k \\ 1 & \text{for } j = k \end{cases},$$

$$W_{jk} = \operatorname{Re}(A_j)\operatorname{Re}(A_k) + \operatorname{Im}(A_j)\operatorname{Im}(A_k) + \operatorname{Re}(B_j)\operatorname{Re}(B_k) + \operatorname{Im}(B_j)\operatorname{Im}(B_k),$$

$$V_{jk} = \operatorname{Re}(A_j)\operatorname{Re}(B_k) + \operatorname{Im}(A_j)\operatorname{Im}(B_k) + \operatorname{Re}(B_j)\operatorname{Re}(A_k) + \operatorname{Im}(B_j)\operatorname{Im}(A_k),$$

$$w_{jkn} = 0, \text{ if } j + k - n \neq 2r, r = 0, 1, 2, \dots, k,$$

$$w_{jkn} = \frac{(\angle j + n - k)(\angle k + n - j)(\angle j + k - n) \left(\angle \frac{j + k + n}{2}\right)^2}{(\angle j + k + n + 1) \left[\left(\angle \frac{j + n - k}{2}\right) \left(\angle \frac{k + n - j}{2}\right) \left(\angle \frac{j + k - n}{2}\right) \right]^2}$$

$$\text{if } j + k - n = 2r, r = 0, 1, 2, \dots, k.$$

$$v_{jkn} = 0, \text{ if } j + k - n \neq 2r + 1, r = 0, 1, 2, \dots, k,$$

and

$$v_{jkn} = \frac{(2n+1)(j+k-n)(j+n-k+1)(k+n-j+1)(\angle j+n-k+1)(\angle k+n-j+1)(\angle j+k-n-1) \left(\angle \frac{j+k+n+1}{2}\right)^2}{4(\angle j+k+n+1) \left[\left(\angle \frac{j+n-k+1}{2}\right) \left(\angle \frac{k+n-j+1}{2}\right) \left(\angle \frac{j+k-n-1}{2}\right) \right]^2}$$

if $j + k - n = 2r + 1, r = 0, 1, 2, \dots, k$

where

Im = Imaginary part,

Re = Real part,

\angle = factorial.

The Legendre polynomials $P_n(\cos \theta)$ were computed by authors (2) up to the high necessary order n by a recurrence relation

$$(n+1) P_{n+1}(x) - (2n+1)x P_n(x) + n P_{n-1}(x) = 0$$

$$\text{starting with } P_0(x) = 1$$

$$P_1(x) = x$$

$$P_n(-x) = (-1)^n P_n(x)$$

For the wanted scattering function

$$4\pi \cdot f(\theta) = 1 + \sum_{n=1}^{\infty} a_n \cdot P_n(\cos \theta)$$

$$\cos \theta = x$$

only the sum of the products $a_n \cdot P_n$ has to be formed.

Since the input of the data a_n and P_n was the only time-consuming work, the symmetry relation

$$P_n(-x) = (-1)^n P_n(x)$$

was used to necessitate the input of P_n for the range of $0^\circ \leq \theta \leq 90^\circ$ only and let the computer fill in the P_n for $90^\circ \leq \theta \leq 180^\circ$ by use of this symmetry relation.

The $f(\theta)$ were computed for scattering angles $\theta = 2^\circ, 5^\circ, 10^\circ, 15^\circ, 20^\circ, 25^\circ, \dots, 160^\circ, 165^\circ, 170^\circ, 175^\circ, 178^\circ$, adjusted to the range of possible scattering measurements, the particle size parameter α took values $\alpha = 1, 2, 3, 4, 5, 6, 8, 10, 15, 20, 25, 30$, according to the values for which the coefficients a_n had been computed in (3). The real index of refraction m covered the range $m = 1.20, 1.25, 1.30, 1.33, 1.44, 1.50, 1.55, 1.60, 2.0, \infty$ for aerosols.

The number of angles θ turned out to be chosen too small in order to represent $f(\theta)$ exactly enough for higher α (15 to 30), because $f(\theta)$ goes through many maxima and minima when scanning $0^\circ \leq \theta \leq 180^\circ$. Since an experimental investigation by Love (5) has shown that $f(\theta)$ probably cannot be truly represented in all maxima and minima (no unique $-\alpha$ mixtures of particles finite beams of radiation), the calculations have not been extended over more intermediate θ - values.

Tables and Graphs

The computer printout gives $4\pi \cdot f(\theta)$ values written in sequence according to the 37 θ - values

$$\begin{array}{lllll}
 \theta_1 = 2^\circ, & \theta_2 = 5^\circ, & \theta_3 = 10^\circ, & \theta_4 = 15^\circ, & \theta_5 = 20^\circ \\
 \theta_6 = 25^\circ & " & " & " & \theta_{10} = 45^\circ \\
 \theta_{11} = 50^\circ & " & " & " & \theta_{15} = 70^\circ \\
 \theta_{16} = 75^\circ & " & " & " & \theta_{20} = 95^\circ \\
 \theta_{21} = 100^\circ & " & " & " & \theta_{25} = 120^\circ \\
 \theta_{26} = 125^\circ & " & " & " & \theta_{30} = 145^\circ \\
 \theta_{31} = 150^\circ & " & " & " & \theta_{35} = 170^\circ \\
 \theta_{36} = 175^\circ, & \theta_{37} = 178^\circ & & &
 \end{array}$$

The $4\pi \cdot f(\theta)$ - values are given to seven decimal places; however 5 are only useful because of the limitations of the tables (2), (3). The last two places may be used for round off.

A limited number of $f(\theta)$ - values was graphed to have an immediate picture of the function. The ordinate in the graphs is signed $f(\theta)$, although the factor 4π is still connected to $f(\theta)$.

The graphs $4\pi f(\theta)$ for α - values 15, 20, 25, 30 do not truly represent the scattering function because the θ - steps are too wide to give enough information over the many maxima and minima of $f(\theta)$, which often extend over a range $\Delta\theta = 2$ to 5° only.

Computer - Program for

$$4\pi f(\theta) = 1 + \sum_{n=1}^N a_n \cdot P_n(\cos \theta)$$

```

/INPUT
/JOB GO
      DIMENSION P(70,37),A(70),F(37)
      READ (5,11) ((P(1,J),I=1,70),J=1,19)
11    FORMAT(10F8.0)
      DO 111 J=1, 18
      DO 111 I=1,69,2
      K=J+19
      L=-J+19
      P(I,K)=-P(I,L)
111   P(I+1,K)=P(I+1,L)
12    READ (5,13)N
13    FORMAT (14)
      IF(N) 14,20,14
14    READ (5,10) (A(I),I=1,N)
10    FORMAT(7F11.0)
      DO 15 J=1,37
15    F(J)=1.0
      DO 16 K=1,37
      DO 16 J=1,N
16    F(K) =F(K)+A(J)*P(J,K)
      WRITE (6,17) F
17    FORMAT (1P5E16.7)
      GO TO 12
20    STOP
      END
/DATA

```


$P(I,J)$ = Legendre polynomials $P_n(\cos \theta)$ for 37 θ - values and orders n up to $N = 70$.


$A(I)$ = Scattering coefficients a_n


$F(K) = 4\pi f(\theta)$


References


1. Mie, G., "Beitrage zur Optik truber Medien", Ann. Physik, 25, 377-445 (1908).
2. Clark, G. C., S. W. Churchill, "Tables of Legendre Polynomials", Univ. of Michigan Press, Engineering Research Inst. Publications, 1957.
3. Chu, C. M., G. C. Clark, S. W. Churchill, "Angular Distribution Coefficients for Radiation Scattered by a Spherical Particle", Journ. Optic. Soc. America, 47, (1957), or in Univ. of Michigan Press, Engineering Research Instit. Publications.
4. Hertel, W., "Zur Theorie der Licht streuung durch trube Schichten", Das Licht, 40, 141-43, 165, 190-91, 214-15, 232-34, (1940)
5. Love, T. J. and R. A. Wheasler, "An Experimental Study of Infra-Red Scattering by Clouds of Particles", Aerospace Research Laboratories, ARL 64-109, June 1964.




















Fig. 8

National Aeronautics and Space Administration
WASHINGTON, D. C.
OFFICIAL BUSINESS

FIRST CLASS MAIL

POSTAGE AND FEES PAID
NATIONAL AERONAUTICS AND
SPACE ADMINISTRATION

070 001 49 01 30S 6-092 00903
AIR FORCE WEAPONS LABORATORY/AFAL/
KIRTLAND AIR FORCE BASE, NEW MEXICO 8711

ALL MISS IADLETT E. CAYOVA, CHIEF TECH
LIBRARY 70117

POSTMASTER: If Undeliverable (Section 15:
Postal Manual) Do Not Return

"The aeronautical and space activities of the United States shall be conducted so as to contribute . . . to the expansion of human knowledge of phenomena in the atmosphere and space. The Administration shall provide for the widest practicable and appropriate dissemination of information concerning its activities and the results thereof."

—NATIONAL AERONAUTICS AND SPACE ACT OF 1958

NASA SCIENTIFIC AND TECHNICAL PUBLICATIONS

TECHNICAL REPORTS: Scientific and technical information considered important, complete, and a lasting contribution to existing knowledge.

TECHNICAL NOTES: Information less broad in scope but nevertheless of importance as a contribution to existing knowledge.

TECHNICAL MEMORANDUMS: Information receiving limited distribution because of preliminary data, security classification, or other reasons.

CONTRACTOR REPORTS: Scientific and technical information generated under a NASA contract or grant and considered an important contribution to existing knowledge.

TECHNICAL TRANSLATIONS: Information published in a foreign language considered to merit NASA distribution in English.

SPECIAL PUBLICATIONS: Information derived from or of value to NASA activities. Publications include conference proceedings, monographs, data compilations, handbooks, sourcebooks, and special bibliographies.

TECHNOLOGY UTILIZATION PUBLICATIONS: Information on technology used by NASA that may be of particular interest in commercial and other non-aerospace applications. Publications include Tech Briefs, Technology Utilization Reports and Notes, and Technology Surveys.

Details on the availability of these publications may be obtained from:

SCIENTIFIC AND TECHNICAL INFORMATION DIVISION
NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

Washington, D.C. 20546